10/626,380 Search Wook 10/11/00

## d his

(FILE 'HOME' ENTERED AT 12:05:47 ON 11 OCT 2006)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, JAPIO' ENTERED AT 12:06:09 ON 11 OCT 2006

	OC1 2006	·				
L1	235	S (ANTI NITROTYROSINE ANTIBOD?)				
L2	1	S L1 AND DNPH?				
L3	69	S L1 AND (OXIDAT?)				
L4		S L3 AND STRESS?				
L5	20	DUPLICATE REMOVE L4 (19 DUPLICATES REMOVED)				
L6	6681 S DINITROPHENYLHYDRAZINE?					
L7	. 2	S L6 AND L5				
L8	2	DUPLICATE REMOVE L7 (0 DUPLICATES REMOVED)				

=>

## d his

(FILE 'HOME' ENTERED AT 12:05:47 ON 11 OCT 2006)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, JAPIO' ENTERED AT 12:06:09 ON 11 OCT 2006

- L1 235 S (ANTI NITROTYROSINE ANTIBOD?)

  L2 1 S L1 AND DNPH?

  L3 69 S L1 AND (OXIDAT?)

  L4 39 S L3 AND STRESS?

  L5 20 DUPLICATE REMOVE L4 (19 DUPLICATES REMOVED)

  L6 6681 S DINITROPHENYLHYDRAZINE?
- L7 2 S L6 AND L5
- L8 2 DUPLICATE REMOVE L7 (0 DUPLICATES REMOVED)

```
ANSWER 2 OF 2 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights
     reserved on STN
     95261594 EMBASE
AN
DN
     1995261594
     Reactive species in ischemic rat lung injury: Contribution of
ΤI
     peroxynitrite.
     Ischiropoulos H.; Al-Mehdi A.B.; Fisher A.B.
ΔII
     Institute for Environmental Medicine, John Morgan Bldg., Univ. of
CS
     Pennsylvania, 3620 Hamilton Walk, Philadelphia, PA 19104-6068, United
     States
     American Journal of Physiology - Lung Cellular and Molecular Physiology,
SO
     (1995) Vol. 269, No. 2 13-2, pp. L158-L164. .
     ISSN: 1040-0605 CODEN: APLPE7
     United States
CY
     Journal; Article
DT
            Physiology
FS
     002
             Drug Literature Index
     037
LA
     English
\mathtt{SL}
     English
     Entered STN: 26 Sep 1995
ED
     Last Updated on STN: 26 Sep 1995
     Lung ischemia-reperfusion represents a potentially important mechanism for
AΒ
     diverse forms of tissue injury associated with decreased pulmonary flow.
     Previous studies demonstrated oxidative injury in
     ischemic-reperfused lungs. The present study was designed to evaluate the
     contribution of nitric oxide and peroxynitrite in tissue injury.
     levels of the stable decomposition products of nitric oxide and
     peroxynitrite, nitrite plus nitrate, were twofold greater then control
     during reperfusion after 60 min of ischemia. Inhibition of nitric oxide
     synthesis by endotracheal insufflation of 5 mM N(G)-nitro-L-arginine
     methyl ester, 30 min before the induction of ischemia, decreased the
     production of lung thiobarbituric acid reactive substances (TBARS) by 67%
     (P < 0.05, n = 5), TBARS released into the lung perfusate by 55% (P <
     0.05, n = 5), lung-conjugated dienes by 61% (P < 0.05, n = 5), and
     dinitrophenylhydrazine-reactive protein carbonyl levels by 86% (P
     < 0.05, n = 5). Amino acid analysis of tissue homogenates from lungs
     exposed to 60 min of ischemia and 60 min of reperfusion revealed a
     1.8-fold (P < 0.05, n = 5) increase in nitrotyrosine concentration
     compared with 2 h continuously perfused lungs. Inhibition of nitric oxide
     synthesis abolished the increase in nitrotyrosine levels. Furthermore,
     lungs exposed to 60 min of reperfusion after 60 min of ischemia showed
     specific binding of an anti-nitrotyrosine
     antibody. In reperfused tissues, antibody binding was observed
     throughout the lung. The binding was blocked with excess of
     nitrotyrosine, and minimal binding was observed in nonperfused blood-free
     control lungs. These results indicate that a strong oxidant derived from
     nitric oxide consistent with the reactivity of peroxynitrite contributes
     to the oxidative injury of isolated rat lung from
     ischemia-reperfusion.
CT
     Medical Descriptors:
     *lung injury: ET, etiology
     *lung perfusion
     animal cell
     animal experiment
     animal tissue
     article
     controlled study
     lung blood flow
     male
     nonhuman
       oxidation
       oxidative stress
     oxygen transport
     pathophysiology
```

priority journal
rat
reperfusion
Drug Descriptors:
\*n(g) nitroarginine methyl ester: PD, pharmacology
\*nitric oxide: EC, endogenous compound
\*oxidizing agent: EC, endogenous compound
\*thiobarbituric acid reactive substance: EC, endogenous compound
peroxynitrite: EC, endogenous compound
RN (n(g) nitroarginine methyl ester) 50903-99-6; (nitric oxide) 10102-43-9

=>

```
ANSWER 6 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN
AN 1975:27408 CAPLUS
DN
     82:27408
ED
     Entered STN: 12 May 1984
     Chemical modification of nucleic acids. II. Reaction
TT
     of calf thymus DNA with hydrazine and 2,4-dinitrophenylhydrazine
     Tsai, Kuang-Hsin; Kantesaria, P.; Marfey, P.
AU
CS
     Dep. Biol. Sci., State Univ. New York, Albany, NY, USA
     Physiological Chemistry and Physics (1974), 6(4), 353-66
so
     CODEN: PLCHB4; ISSN: 0031-9325
DT
     Journal
LΑ
     English
CC
     6-2 (General Biochemistry)
     2,4-Dinitrophenylhydrazine (DNPH) reacted with deoxyadenosine,
     deoxyguanosine, and deoxycytidine under mild conditions, but not with
     thymidine. Only deoxycytidine reacted with hydrazine at pH 6. Treatment
     of DNA with DNPH at pH 4 led to the incorporation of 1
     DNPH group/111-165 deoxynucleotide residues. Treatment of DNA 1st
     with hydrazine at pH 6 followed by treatment with excess
     1-fluoro-2,4-dinitrobenzene at pH 8.2 afforded derivs. in which, depending
     on exptl. conditions, 1 DNPH group was introduced/29-528
     deoxynucleotide residues. The derivs. obtained exhibited high mol. weight
     and retained the native structure. The covalently attached DNPH
     chromophore in DNA may be a useful absorption probe in a study of its
     interaction with other mols. or ions.
     DNA reaction hydrazine dinitrophenylhydrazine
ST
     Deoxyribonucleic acids
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (reaction of, with hydrazine and dinitrophenylhydrazine)
IT
     119-26-6
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (reaction of, with DNA)
IT
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (reaction of, with DNA hydrazine derivative)
IT
     961-07-9
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (reaction of, with dinitrophenylhydrazine)
IT
     951-77-9
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (reaction of, with dinitrophenylhydrazine and hydrazine)
IT
     302-01-2, reactions
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (with DNA)
IT
     58-61-7, reactions
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (with dinitrophenylhydrazine)
```

```
ANSWER 6 OF 7 CAPLUS COPYRIGHT 2006 ACS on STN
     1975:27408 CAPLUS
AN
DN
     82:27408
     Entered STN: 12 May 1984
ED
     Chemical modification of nucleic acids. II. Reaction
TI
     of calf thymus DNA with hydrazine and 2,4-dinitrophenylhydrazine
     Tsai, Kuang-Hsin; Kantesaria, P.; Marfey, P.
ΑU
     Dep. Biol. Sci., State Univ. New York, Albany, NY, USA
CS
SO
     Physiological Chemistry and Physics (1974), 6(4), 353-66
     CODEN: PLCHB4; ISSN: 0031-9325
DT
     Journal
     English
LA
CC
     6-2 (General Biochemistry)
     2,4-Dinitrophenylhydrazine (DNPH) reacted with deoxyadenosine,
AΒ
     deoxyguanosine, and deoxycytidine under mild conditions, but not with
     thymidine. Only deoxycytidine reacted with hydrazine at pH 6. Treatment
     of DNA with DNPH at pH 4 led to the incorporation of 1
     DNPH group/111-165 deoxynucleotide residues. Treatment of DNA 1st
     with hydrazine at pH 6 followed by treatment with excess
     1-fluoro-2,4-dinitrobenzene at pH 8.2 afforded derivs. in which, depending
     on exptl. conditions, 1 DNPH group was introduced/29-528
     deoxynucleotide residues. The derivs. obtained exhibited high mol. weight
     and retained the native structure. The covalently attached DNPH
     chromophore in DNA may be a useful absorption probe in a study of its
     interaction with other mols. or ions.
     DNA reaction hydrazine dinitrophenylhydrazine
ST
     Deoxyribonucleic acids
IT
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (reaction of, with hydrazine and dinitrophenylhydrazine)
IT
     119-26-6
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (reaction of, with DNA)
IT
     70-34-8
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (reaction of, with DNA hydrazine derivative)
IT
     961-07-9
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (reaction of, with dinitrophenylhydrazine)
IT
     951-77-9
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (reaction of, with dinitrophenylhydrazine and hydrazine)
IT
     302-01-2, reactions
     RL: RCT (Reactant); RACT (Reactant or reagent)
        (with DNA)
IT
     58-61-7, reactions
     RL: RCT (Reactant); RACT (Reactant or reagent)
```

(with dinitrophenylhydrazine)

ANSWER 1 OF 226 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

AN 2001:462605 BIOSIS

DN PREV200100462605

TI Processes for detecting polynucleotides, determining genetic mutations or defects in genetic material, separating or isolating nucleic acid of interest from samples, and useful compositions of matter and multihybrid complex compositions.

AU Engelhardt, Dean L. [Inventor, Reprint author]; Rabbani, Elazar [Inventor]

CS New York, NY, USA
ASSIGNEE: Enzo Diagnostics, Inc., New York, NY, USA; c/o Enzo Biochem,
Inc., New York, NY, USA

PI US 6221581 20010424

SO Official Gazette of the United States Patent and Trademark Office Patents, (Apr. 24, 2001) Vol. 1245, No. 4. e-file.

CODEN: OGUPE7. ISSN: 0098-1133.

DT Patent

LA English

ED Entered STN: 3 Oct 2001 Last Updated on STN: 22 Feb 2002

Double hybrid or multihybrid probes and compositions are usefully combined with capture assay and immobilization to provide for detection processes in which target polynucleotides can be detected or the presence or absence of genetic mutations or defects in genetic material can be determined. The capture assay involves capturing a hybrid structure, e.g., single hybrid, double hybrid or multihybrid, or capturing a complex formed by reacting a hybrid structure with a complex forming moiety, e.g., protein, such as a binding protein including an antibody. Immobilization can also be employed prior to hybridization or complexation in which instance a polynucleotide probe can be fixed to a matrix or solid support, e.g., natural or synthetic. Capture and immobilization can be carried out using direct and indirect binding and attachment techniques. Targets can be detected directly or indirectly by using a signal generating moiety and labels.

NCL 435006000

CC General biology - Miscellaneous 00532

IT Major Concepts

Molecular Genetics (Biochemistry and Molecular Biophysics); Methods and Techniques

IT Methods & Equipment

genetic material defect detection: detection method; genetic mutation determination: determination method; nucleic acid isolation: isolation method; nucleic acid separation: separation method; polynucleotide detection: detection method

IT Miscellaneous Descriptors
 double hybrid probes; multihybrid probes

L7 ANSWER 2 OF 226 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation

```
ANSWER 3 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN
AN
     2000:633957 CAPLUS
DN
     134:82839
     Entered STN: 13 Sep 2000
ED
     Immobilization of protein monolayers on planar solid
TI
     supports
ΑU
     Dubrovsky, Timothy B.
     Roche Diagnostic Systems, Inc., Somerville, NJ, USA
CS
     Protein Architecture (2000), 25-54. Editor(s): Lvov, Yuri;
SO
     Moehwald, Helmuth. Publisher: Marcel Dekker, Inc., New York, N. Y.
     CODEN: 69AHGU
DT
     Conference; General Review
     English
LA
     9-0 (Biochemical Methods)
CC
     A review with 111 refs. is presented regarding the critical steps
AΒ
     of preparation, activation, and characterization of self-assembled monolayers
     of silane mols: The strategies for synthesis of two self-assembling
     systems that can be used for covalent immobilization of protein monolayers
     on solid supports are described. The first class of
     self-assembling system is alkylsilane compds. with different terminal
     functionalities. The general procedures for silanization of silicon and
     glass surfaces as well as the anal. methods for characterization of
     self-assembled monolayers are described. Characterization by
     spectroscopies, ellipsometry, and contact angles verified that synthetic
     routes employed lead to well-defined surfaces with controlled mol.
     architecture that can be routinely used in various biomaterials
     investigations. It has been demonstrated that monolayers of mitochondrial
     cytochrome P450scc and oriented antibody layers can be
     transferred from the air-water interface to the silanized quartz supports
     without damage to the structure. The second class of self-assembling
     system discussed is \omega-substituted alkanethiols chemisorbed onto the
     surface of gold. Alkanethiol monolayers are stable, permit the
     introduction of a variety of functional groups onto surfaces, and can be
     well organized. These monolayers can host either active groups or
     affinity ligands for the specific binding of protein mols. This approach
     to synthesis of model surfaces may find use in diagnostic assays
     and affinity chromatog.
     review protein immobilization planar support
st
IT
     Proteins, general, reactions
     RL: PEP (Physical, engineering or chemical process); RCT (Reactant); PROC
     (Process); RACT (Reactant or reagent)
        (immobilization of protein monolayers on planar solid
        supports)
IT
     Immobilization, biochemical
        (protein; immobilization of protein monolayers on planar solid
              THERE ARE 111 CITED REFERENCES AVAILABLE FOR THIS RECORD
RE.CNT
        111
RE
(1) Ahluvalia, A; Biosens Bioelectron 1991, V6, P133
(2) Ahluvalia, A; Biosens Bioelectron 1992, V7, P207
(3) Amador, S; Langmuir 1993, V9, P812 CAPLUS
(4) Anon; Chemical Sensors and Microinstrumentation 1989
(5) Anon; Practical Protein Chemistry 1986, P386
(6) Bain, C; J Am Chem Soc 1989, V111, P321 CAPLUS
(7) Balachander, N; Langmuir 1990, V6, P1621 CAPLUS
(8) Beyer, D; Langmuir 1996, V12, P2514 CAPLUS
(9) Bhatia, S; Anal Biochem 1989, V178, P408 CAPLUS
(10) Bierbaum, K; Langmuir 1995, V11, P512 CAPLUS
(11) Bierbaum, K; Langmuir 1995, V11, P512 CAPLUS
(12) Blankenburg, R; Biochemistry 1989, V28, P8214 CAPLUS
(13) Blodgett, K; Phys Rev 1937, V51, P964 CAPLUS
(14) Bruckenstein, S; Anal Chem 1994, V66, P1847 CAPLUS
(15) Cohen, S; J Chem Phys 1986, V90, P3054 CAPLUS
(16) Darst, S; Biophys J 1991, V59, P387 CAPLUS
```

```
(17) Dubrovsky, T; Anal Chem 1999, V71, P327 CAPLUS
```

- (18) Dubrovsky, T; Biosens Bioelectron 1993, V8, P377 CAPLUS
- (19) Dubrovsky, T; Langmuir in press
- (20) Dubrovsky, T; Russian Chem Bull 1993, V42, P1488
- (21) Dubrovsky, T; Sensors and Actuators B 1994, V22, P69
- (22) Dubrovsky, T; Thin Solid Films 1996, V284, P698
- (23) Dubrovsky, T; unpublished results 1994
- (24) Dunaway, D; Langmuir 1994, V10, P3598 CAPLUS
- (25) Edmiston, P; J Am Chem Soc 1997, V119, P560 CAPLUS
- (26) Erokhin, V; Biosens Bioelectron 1995, V10, P25 CAPLUS
- (27) Erokhin, V; Studia Biophys 1989, V132, P97 CAPLUS
- (28) Facci, P; Thin Solid Films 1994, V243, P403 CAPLUS
- (29) Fang, J; Langmuir 1995, V11, P4366 CAPLUS
- (30) Fatunmbi, H; Anal Chem 1993, V65, P2048 CAPLUS
- (31) Gupta, V; Langmuir 1996, V12, P2587 CAPLUS
- (32) Gupta, V; Science 1998, V279, P2077 CAPLUS
- (33) Guryev, O; Biochem Mol Biol Int 1996, V39, P205 CAPLUS
- (34) Guryev, O; Langmuir 1997, V13, P299 CAPLUS
- (35) Haller, I; J Am Chem Soc 1978, V100, P8050 CAPLUS
- (36) Harder, P; Langmuir 1997, V13, P445 CAPLUS
- (37) Haussling, L; Angew Chem Int Ed 1991, V30, P569
- (38) Haussling, L; Langmuir 1991, V7, P1837
- (39) Haussling, L; Langmuir 1991, V7, P1837
- (40) Heise, A; Langmuir 1997, V13, P723 CAPLUS
- (41) Hemmila, I; Anal Biochem 1984, V137, P335 CAPLUS
- (42) Hermanson, G; Immobilized Affinity Ligand Techniques 1992, P103
- (43) Herron, J; Langmuir 1992, V8, P1413 CAPLUS
- (44) Hoffmann, P; Langmuir 1997, V13, P1877 CAPLUS
- (45) Hong, H; Anal Chem 1993, V65, P1635 CAPLUS
- (46) Hong, H; Langmuir 1994, V10, P153 CAPLUS
- (47) Jonsson, U; Thin Solid Films 1985, V124, P117
- (48) Kallury, K; Langmuir 1994, V10, P492 CAPLUS
- (49) Karpovich, D; Langmuir 1994, V10, P3315 CAPLUS (50) Katz, E; J Electroanal Chem 1990, V291, P257
- (51) Knichel, M; Sensors and Actuators B 1995, V28, P85
- (52) Knoll, W; Int Patent Appl WO 92/10757
- (53) Kohler, J; J Chromatogr 1986, V352, P275
- (54) Kurth, D; Angew Chem Int Ed 1992, V31, P336
- (55) Kurth, D; Langmuir 1993, V9, P2965 CAPLUS
- (56) Kurth, D; Langmuir 1995, V11, P3061 CAPLUS
- (57) Langmuir, I; J Am Chem Soc 1938, V60, P1351 CAPLUS
- (58) Leckband, D; Biochemistry 1995, V34, P11467 CAPLUS
- (59) Lehmann, A; FEBS Lett 1984, V176, P360 CAPLUS
- (60) Lion-Dagan, M; J Chem Soc Chem Commun V1994, P2741
- (61) Lotzbeyer, T; Sensors and Actuators B 1996, V33, P50
- (62) Lu, B; Thin Solid Films 1994, V240, P138 CAPLUS
- (63) Lvov, Y; Biol Membr 1991, V9, P1477
- (64) Lyn, M; Immobilized Enzymes, Antigens, Antibodies, and Peptides 1978
- (65) Malmqvist, M; US 4833093 1989 CAPLUS
- (66) Maoz, R; J Colloid Interface Sci 1984, V100, P465 CAPLUS
- (67) Moon, J; Langmuir 1996, V12, P4621 CAPLUS
- (68) Mrksich, M; J Am Chem Soc 1995, V117, P12009 CAPLUS
- (69) Nakano, K; Anal Sci 1993, V9, P133 CAPLUS
- (70) Netzer, L; Thin Solid Films 1983, V100, P67 CAPLUS
- (71) Nevernov, I; Thin Solid Films 1994, V252, P1 CAPLUS
- (72) Nicolini, C; Biochim Biophys Acta 1993, V1158, P273 CAPLUS
- (73) Paddeu, S; Thin Solid Films 1995, V268, P108 CAPLUS
- (74) Pale-Grosdemange, C; J Am Chem Soc 1991, V113, P12 CAPLUS
- (75) Parikh, A; J Phys Chem 1994, V98, P7577 CAPLUS
- (76) Pomerantz, M; Thin Solid Films 1985, V132, P153 CAPLUS
- (77) Prime, K; J Am Chem Soc 1993, V115, P10714 CAPLUS
- (78) Prime, K; Science 1991, V252, P1164 CAPLUS
- (79) Reiter, R; Langmuir 1993, V9, P2430 CAPLUS
- (80) Roberts, G; Langmuir-Blodgett Films 1990

- (81) Sagiv, J; J Am Chem Soc 1980, V102, P92 CAPLUS
- (82) Salesse, C; Biochemistry 1990, V29, P4567 CAPLUS
- (83) Spinke, J; J Chem Phys 1993, V99, P7012 CAPLUS
- (84) Spinke, J; Langmuir 1993, V9, P1821 CAPLUS
- (85) Spinke, J; Langmuir 1993, V9, P1821 CAPLUS
- (86) Staros, J; Anal Biochem 1986, V156, P220 CAPLUS
- (87) Stayton, P; J Am Chem Soc 1992, V114, P9298 CAPLUS
- (88) Tidswell, I; Phys Rev B 1990, V41, P1111 CAPLUS
- (89) Tillman, N; J Am Chem Soc 1988, V111, P6136
- (90) Tronin, A; Langmuir 1995, V11, P385 CAPLUS
- (91) Tronin, A; Langmuir 1996, V12, P3272 CAPLUS
- (92) Troughton, E; Langmuir 1988, V4, P365 CAPLUS
- (93) Turko, I; Thin Solid Films 1991, V205, P113 CAPLUS
- (94) Turko, I; Thin Solid Films 1992, V210, P710
- (95) Ulman, A; Adv Mater 1990, V2, P573 CAPLUS
- (96) Ulman, A; An Introduction to Ultrathin Organic Films: From Langmuir-Blodgett to Self-Assembly 1991
- (97) Ulman, A; An Introduction to Ultrathin Organic Films: From Langmuir-Blodgett to Self-Assembly 1991
- (98) Uzgiris, E; Biochem J 1987, V242, P293 CAPLUS
- (99) Uzgiris, E; Nature 1983, V301, P125 CAPLUS
- (100) Wachter, E; FEBS Lett 1973, V35, P97 CAPLUS
- (101) Wasserman, S; J Am Chem Soc 1989, V111, P5852 CAPLUS
- (102) Wasserman, S; Langmuir 1989, V5, P1074 CAPLUS
- (103) Welsch, T; J Chromatogr 1990, V506, P97 CAPLUS
- (104) Wilchek, M; Avidin-Biotin Technology V184(13), P123
- (105) Willner, I; J Am Chem Soc 1992, V114, P10965 CAPLUS
- (106) Willner, I; J Am Chem Soc 1993, V115, P4937 CAPLUS
- (107) Wirth, M; Science 1997, V275, P44 CAPLUS
- (108) Wood, L; J Am Chem Soc 1997, V119, P571 CAPLUS
- (109) Wood, L; J Am Chem Soc 1997, V119, P571 CAPLUS
- (110) Xia, Y; J Am Chem Soc 1995, V117, P9576 CAPLUS
- (111) Yamaki, M; Langmuir 1993, V9, P3154 CAPLUS

```
ANSWER 11 OF 13 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights
     reserved on STN
AN
     1999358516 EMBASE
     Microscale determinations using solid phase assays: Applications
ΤI
     to biochemical, clinical and biotechnological sectors. A review.
ΑU
     Vynios D.H.
     D.H. Vynios, Laboratory of Biochemistry, Section of Organic Chemistry,
CS
     Department of Chemistry, 261 10 Patras, Greece
     Journal of Liquid Chromatography and Related Technologies, (1999
SO
     ) Vol. 22, No. 17, pp. 2555-2574. .
     Refs: 36
     ISSN: 1082-6076 CODEN: JLCTFC
     United States
CY
DT
     Journal; General Review
          Biophysics, Bioengineering and Medical Instrumentation
FS
             Clinical Biochemistry
     English
LA
     English
SL
     Entered STN: 29 Oct 1999
ED
     Last Updated on STN: 29 Oct 1999
     The assays that have one of the reactant species immobilized
AB
     onto a solid support are described as solid phase
     assays. During the last 20 years a large number of such
     assays has been developed, the majority of which are quantitative
     analytical methods known under the general term ELISA (Enzyme Linked
     ImmunoSorbent Assay). Solid phase assays, in general,
     have widely been used in Biochemistry, Clinical Chemistry, and
     Biotechnology, mainly for analytical purposes, and for the detection of
     specific macromolecules or the study of interactions between various
     molecules, as well.
CT
    Medical Descriptors:
       *assay
       enzyme linked immunosorbent assay
     clinical chemistry
     biotechnology
     polymerase chain reaction
     zymography
     human
       review
     Drug Descriptors:
     *messenger RNA: EC, endogenous compound
     *DNA: EC, endogenous compound
     *streptavidin
     *metalloproteinase: EC, endogenous compound
     *hyaluronic acid
     *hyaluronidase: EC, endogenous compound
     *autoantibody: EC, endogenous compound
     *oligosaccharide
     *lysozyme
     *aggrecan
     polystyrene
       monoclonal antibody
     antigen
     biotin
     avidin
     protein
     proteoglycan
     glycosaminoglycan
     1 (3 dimethylaminopropyl) 3 ethylcarbodiimide
     glucose: EC, endogenous compound
     oxidoreductase
     peroxidase
     peroxide
     bilirubin: EC, endogenous compound
```

nitrite: EC, endogenous compound
ketone body: EC, endogenous compound
urobilinogen: EC, endogenous compound
chorionic gonadotropin: EC, endogenous compound
collagen

RN (DNA) 9007-49-2; (streptavidin) 9013-20-1; (metalloproteinase) 81669-70-7;
(hyaluronic acid) 31799-91-4, 9004-61-9, 9067-32-7; (hyaluronidase)
9001-54-1, 9055-18-9; (lysozyme) 9001-63-2; (polystyrene) 9003-53-6;
(biotin) 58-85-5; (protein) 67254-75-5; (1 (3 dimethylaminopropyl) 3
ethylcarbodiimide) 1892-57-5, 25952-53-8, 7084-11-9; (glucose) 50-99-7,
84778-64-3; (oxidoreductase) 9035-73-8, 9035-82-9, 9037-80-3, 9055-15-6;
(peroxidase) 9003-99-0; (peroxide) 14915-07-2; (bilirubin) 18422-02-1,
635-65-4; (nitrite) 14797-65-0; (urobilinogen) 11000-27-4; (chorionic gonadotropin) 9002-61-3; (collagen) 9007-34-5

```
ANSWER 11 OF 13 EMBASE COPYRIGHT (c) 2006 Elsevier B.V. All rights
     reserved on STN
     1999358516 EMBASE
AN
     Microscale determinations using solid phase assays: Applications
TI
     to biochemical, clinical and biotechnological sectors. A review.
ΑU
     Vynios D.H.
     D.H. Vynios, Laboratory of Biochemistry, Section of Organic Chemistry,
CS
     Department of Chemistry, 261 10 Patras, Greece
     Journal of Liquid Chromatography and Related Technologies, (1999
SO
     ) Vol. 22, No. 17, pp. 2555-2574. .
     Refs: 36
     ISSN: 1082-6076 CODEN: JLCTFC
     United States
CY
     Journal; General Review
DT
             Biophysics, Bioengineering and Medical Instrumentation
FS
     027
             Clinical Biochemistry
     029
LA
     English
    English
SL
     Entered STN: 29 Oct 1999
ED
     Last Updated on STN: 29 Oct 1999
     The assays that have one of the reactant species immobilized
AB
     onto a solid support are described as solid phase
     assays. During the last 20 years a large number of such
     assays has been developed, the majority of which are quantitative
     analytical methods known under the general term ELISA (Enzyme Linked
     ImmunoSorbent Assay). Solid phase assays, in general,
     have widely been used in Biochemistry, Clinical Chemistry, and
     Biotechnology, mainly for analytical purposes, and for the detection of
     specific macromolecules or the study of interactions between various
     molecules, as well.
CT
     Medical Descriptors:
       *assay
       enzyme linked immunosorbent assay
     clinical chemistry
     biotechnology
     polymerase chain reaction
     zymography
     human
       review
     Drug Descriptors:
     *messenger RNA: EC, endogenous compound
     *DNA: EC, endogenous compound
     *streptavidin
     *metalloproteinase: EC, endogenous compound
     *hyaluronic acid
     *hyaluronidase: EC, endogenous compound
     *autoantibody: EC, endogenous compound
     *oligosaccharide
     *lysozyme
     *aggrecan
     polystyrene
       monoclonal antibody
     antigen
     biotin
     avidin
     protein
     proteoglycan
     glycosaminoglycan
     1 (3 dimethylaminopropyl) 3 ethylcarbodiimide
     glucose: EC, endogenous compound
     oxidoreductase
     peroxidase
     peroxide
     bilirubin: EC, endogenous compound
```

ketone body: EC, endogenous compound
urobilinogen: EC, endogenous compound
chorionic gonadotropin: EC, endogenous compound
collagen
RN (DNA) 9007-49-2; (streptavidin) 9013-20-1; (metalloproteinase) 81669-70-7;
(hyaluronic acid) 31799-91-4, 9004-61-9, 9067-32-7; (hyaluronidase)
9001-54-1, 9055-18-9; (lysozyme) 9001-63-2; (polystyrene) 9003-53-6;
(biotin) 58-85-5; (protein) 67254-75-5; (1 (3 dimethylaminopropyl) 3
ethylcarbodiimide) 1892-57-5, 25952-53-8, 7084-11-9; (glucose) 50-99-7,
84778-64-3; (oxidoreductase) 9035-73-8, 9035-82-9, 9037-80-3, 9055-15-6;
(peroxidase) 9003-99-0; (peroxide) 14915-07-2; (bilirubin) 18422-02-1,
635-65-4; (nitrite) 14797-65-0; (urobilinogen) 11000-27-4; (chorionic gonadotropin) 9002-61-3; (collagen) 9007-34-5

nitrite: EC, endogenous compound

PALM Intranet

Application Number

Submit

IDS Flag Clearance for Application 10626380



Content	Mailroom Date	Entry Number	IDS Review	Last Modified	Reviewer
M844	2003-08-21	13	Y 🗹	2006-10 <b>-</b> 11 13:38:18.0	LCook
Update					